

5

Literature Cited

- Ammann, A.P. and A. Lindley Stone 1991. Method for the Comparative Evaluation of Nontidal Wetlands in New Hampshire. Published by the NH Department of Environmental Services. NHDES-WRD-1991-3.
- Apfelbaum, S.I. and C.E. Sams. 1987. Ecology and control of reed canarygrass (*Phalaris arundinacea* L.). *Natural Areas Journal* 7: 69-74.
- Cowardin, L.M., V. Carter, F.C. Golet and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. US Fish and Wildlife Biological Services Program. FWS/OBS-79/31, Washington, DC. 103 pp.
- Durkis, T. 2003. Biological Control of Purple Loosestrife, 1997-2001. New Hampshire Department of Agriculture, Markets & Food, Plant Industry and the New Hampshire Department of Transportation. FHWA-NH-RD-12323Q. 59 pp +
- Golet, F.C., A.J.K. Calhoun, W.R. DeRagon, D.J. Lowry, and A.J. Gold. 1993. Ecology of Red Maple Swamps in the Glaciated Northeast: A Community Profile. Biological Report 12. US Dept. of Interior, Fish and Wildlife Services, Washington, D.C., 151 pp.
- Justice, D., A.K. Deely, and F. Rubin. 2002. New Hampshire Land Cover Assessment: Final Report. Complex Systems Research Center, University of New Hampshire, Durham, NH, 42 p.
- Michigan Department of Environmental Quality. 2008. A Guide to the Control and Management of Invasive Phragmites. Office of the Great Lakes, Aquatic Invasive Species Program, 37 pp.